

Module 2: Set Theory

Operational math bootcamp



Statistical Sciences
UNIVERSITY OF TORONTO

Ichiro Hashimoto

University of Toronto

July 9, 2024

Outline

- Review of basic set theory
- Ordered Sets
- Functions

Introduction to Set Theory

- We define a *set* to be a collection of mathematical objects.
- If S is a set and x is one of the objects in the set, we say x is an element of S and denote it by $x \in S$.
- The set of no elements is called empty set and is denoted by \emptyset .

Definition (Subsets, Union, Intersection)

Let S, T be sets.

- We say that S is a *subset* of T , denoted $S \subseteq T$, if $s \in S$ implies $s \in T$.
- We say that $S = T$ if $S \subseteq T$ and $T \subseteq S$.
- We define the *union* of S and T , denoted $S \cup T$, as all the elements that are in *either* S or T .
- We define the *intersection* of S and T , denoted $S \cap T$, as all the elements that are in *both* S and T .
- We say that S and T are *disjoint* if $S \cap T = \emptyset$.

Some examples

Example

$$\mathbb{N} \subseteq \mathbb{N}_0 \subseteq \mathbb{Z} \subseteq \mathbb{Q} \subseteq \mathbb{R} \subseteq \mathbb{C}$$

Example

Let $a, b \in \mathbb{R}$ such that $a < b$.

Open interval: $(a, b) := \{x \in \mathbb{R} : a < x < b\}$ (a, b may be $-\infty$ or $+\infty$)

Closed interval: $[a, b] := \{x \in \mathbb{R} : a \leq x \leq b\}$

We can also define half-open intervals.

Example

Let $A = \{x \in \mathbb{N} : 3|x\}$ and $B = \{x \in \mathbb{N} : 6|x\}$. Show that $B \subseteq A$.

Proof.

Difference of sets

Definition

Let $A, B \subseteq X$. We define the *set-theoretic difference* of A and B , denoted $A \setminus B$ (sometimes $A - B$) as the elements of X that are in A but *not* in B .

The complement of a set $A \subseteq X$ is the set $A^c := X \setminus A$.

Example

Let $X \subseteq \mathbb{R}$ be defined as $X = \{x \in \mathbb{R} : 0 < x \leq 40\} = (0, 40]$. Then

$X^c =$

Recall that for sets S, T :

- the *union* of S and T , denoted $S \cup T$, is all the elements that are in *either* S and T
- and the *intersection* of S and T , denoted $S \cap T$, is all the elements that are in *both* S and T .

We extend the definition of union and intersection to an arbitrary family of sets as follows:

Definition

Let $S_\alpha, \alpha \in A$, be a family of sets. A is called the *index set*. We define

$$\bigcup_{\alpha \in A} S_\alpha := \{x : \exists \alpha \text{ such that } x \in S_\alpha\},$$

$$\bigcap_{\alpha \in A} S_\alpha := \{x : x \in S_\alpha \text{ for all } \alpha \in A\}.$$

Example

$$\bigcup_{n=1}^{\infty} [-n, n] =$$

$$\bigcap_{n=1}^{\infty} \left(-\frac{1}{n}, \frac{1}{n} \right) =$$

Theorem (De Morgan's Laws)

Let $\{S_\alpha\}_{\alpha \in A}$ be an arbitrary collection of sets. Then

$$\left(\bigcup_{\alpha \in A} S_\alpha \right)^c = \bigcap_{\alpha \in A} S_\alpha^c \quad \text{and} \quad \left(\bigcap_{\alpha \in A} S_\alpha \right)^c = \bigcup_{\alpha \in A} S_\alpha^c$$

Proof.

Since a set is itself a mathematical object, a set can itself contain sets.

Definition

The power set $\mathcal{P}(S)$ of a set S is the set of all subsets of S .

Example

Let $S = \{a, b, c\}$.

Then $\mathcal{P}(S) =$

Another way of building a new set from two old ones is the Cartesian product of two sets.

Definition

Let S, T be sets. The *Cartesian product* $S \times T$ is defined as the set of tuples with elements from S, T , i.e

$$S \times T = \{(s, t) : s \in S \text{ and } t \in T\}.$$

This can also be extended inductively to a finite family of sets.

Ordered set

Definition

A *relation* R on a set X is a subset of $X \times X$. A relation \leq is called a *partial order* on X if it satisfies

- 1 reflexivity:
- 2 transitivity:
- 3 anti-symmetry:

The pair (X, \leq) is called a *partially ordered set*.

A *chain* or *totally ordered set* $C \subseteq X$ is a subset with the property $x \leq y$ or $y \leq x$ for any $x, y \in C$.

Example

The real numbers with the usual ordering, (\mathbb{R}, \leq) are totally ordered.

Example

The power set of a set X with the ordering given by \subseteq , $(\mathcal{P}(X), \subseteq)$ is a partially ordered set.

Example

Let $X = \{a, b, c, d\}$. What is $\mathcal{P}(X)$? Find a chain in $\mathcal{P}(X)$.

$$\mathcal{P}(X) = \{\emptyset, \{a\}, \{b\}, \{c\}, \{d\}, \{a, b\}, \{b, c\}, \{c, d\}, \{b, d\}, \{a, c\}, \{a, d\}, \{a, b, c\}, \{b, c, d\}, \{a, b, d\}, \{a, c, d\}, X\}$$

Example

Consider the set $C([0, 1], \mathbb{R}) := \{f: [0, 1] \rightarrow \mathbb{R} : f \text{ is continuous}\}$.

For two functions $f, g \in C([0, 1], \mathbb{R})$, we define the ordering as $f \leq g$ if $f(x) \leq g(x)$ for $x \in [0, 1]$. Then $(C([0, 1], \mathbb{R}), \leq)$ is a partially ordered set.

Can you think of a chain that is a subset of $(C([0, 1], \mathbb{R}))$?

Definition

A non-empty partially ordered set (X, \leq) is *well-ordered* if every non-empty subset $A \subseteq X$ has a minimum element.

Example:

(\mathbb{N}, \leq) is...

(\mathbb{R}, \leq) is...

Definition

Let (X, \leq) be a partially ordered set and $S \subseteq X$.

Then $x \in X$ is an *upper bound* for S if for all $s \in S$ we have $s \leq x$.

Similarly, $y \in X$ is a *lower bound* for S if for all $s \in S$, $y \leq s$.

If there exists an upper bound for S , we call S *bounded above* and if there exists a lower bound for S , we call S *bounded below*. If S is bounded above and bounded below, we say S is *bounded*.

We can also ask if there exists a least upper bound or a greatest lower bound.

Definition

Let (X, \leq) be a partially ordered set and $S \subseteq X$.

We call $x \in X$ *least upper bound* or *supremum*, denoted $x = \sup S$, if x is an upper bound and for any other upper bound $y \in X$ of S we have $x \leq y$.

Likewise, $x \in X$ is the *greatest lower bound* or *infimum* for S , denoted $x = \inf S$, if it is a lower bound and for any other lower bound $y \in X$, $y \leq x$.

Note that the supremum and infimum of a bounded set do not necessarily need to exist. However, if they do exist they are unique, which justifies the article *the* (exercise). Nevertheless, the reals have a remarkable property, which we will take as an axiom.

Completeness Axiom

Let $S \subseteq \mathbb{R}$ be bounded above. Then there exists $r \in \mathbb{R}$ such that $r = \sup S$, i.e. S has a least upper bound.

By setting $S' = -S := \{-s : s \in S\}$ and noting $\inf S = -\sup S'$, we obtain a similar statement for infima if S is bounded below. As mentioned above, this property is fairly special, for example it fails for the rationals.

Example

Let $S = \{q \in \mathbb{Q} : q^2 < 7\}$. Then S is bounded above in \mathbb{Q} , but there exists no least upper bound in \mathbb{Q} .

There is a nice alternative characterization for suprema in the real numbers.

Proposition

Let $S \subseteq \mathbb{R}$ be bounded above. Then $r = \sup S$ if and only if r is an upper bound and for all $\epsilon > 0$ there exists an $s \in S$ such that $r - \epsilon < s$.

Proof. (\Rightarrow)

Proof. (\Leftarrow)

Using the same trick, we may obtain a similar result for infima.

Example

Consider $S = \{1/n : n \in \mathbb{N}\}$. Then $\sup S = 1$ and $\inf S = 0$.

Functions

Definition

A function f from a set X to a set Y is a subset of $X \times Y$ with the properties:

- 1 For every $x \in X$, there exists a $y \in Y$ such that $(x, y) \in f$
- 2 If $(x, y) \in f$ and $(x, z) \in f$, then $y = z$.

X is called the *domain* of f .

How does this connect to other descriptions of functions you may have seen?

Example

For a set X , the identity function is:

$$1_X : X \rightarrow X, \quad x \mapsto x$$

Definition (Image and pre-image)

Let $f: X \rightarrow Y$ and $A \subseteq X$ and $B \subseteq Y$.

- The *image* of f is the set $f(A) := \{f(x) : x \in A\}$.
- The *pre-image* of f is the set $f^{-1}(B) := \{x : f(x) \in B\}$.

Helpful way to think about it for proofs:

Image: If $y \in f(A)$, then $y \in Y$, and there exists an $x \in A$ such that $y = f(x)$.

Pre-image: If $x \in f^{-1}(B)$, then $x \in X$ and $f(x) \in B$.

Definition (Surjective, injective and bijective)

Let $f: X \rightarrow Y$, where X and Y are sets. Then

- f is *injective* if $x_1 \neq x_2$ implies $f(x_1) \neq f(x_2)$
- f is *surjective* if for every $y \in Y$, there exists an $x \in X$ such that $y = f(x)$
- f is *bijective* if it is both injective and surjective

Example

Let $f: X \rightarrow Y, x \mapsto x^2$.

f is surjective if

f is injective if

f is bijective if

f is neither surjective nor injective if

References

Marcoux, Laurent W. (2019). *PMATH 351 Notes*. url:
<https://www.math.uwaterloo.ca/lwmarcou/notes/pmath351.pdf>

Runde ,Volker (2005). *A Taste of Topology*. Universitext. url:
<https://link.springer.com/book/10.1007/0-387-28387-0>

Zwiernik, Piotr (2022). *Lecture notes in Mathematics for Economics and Statistics*.
url: <http://84.89.132.1/piotr/docs/RealAnalysisNotes.pdf>