

# Module 8: Linear Algebra II

## Operational math bootcamp



Statistical Sciences  
UNIVERSITY OF TORONTO

Ichiro Hashimoto

University of Toronto

July 22, 2024

# Outline

Last time:

- Vector spaces and subspaces
- Linear independence and bases
- Linear maps, null space, range

Today:

- Inverses of linear maps
- Matrices as linear maps
- Determinants
- Inner product spaces

# Recall

## Theorem

Suppose  $\mathbf{u}_1, \dots, \mathbf{u}_n$  is a basis for  $U$  and  $\mathbf{v}_1, \dots, \mathbf{v}_n$  is a basis for  $V$ . Then there exists a unique linear map  $T: U \rightarrow V$  such that  $T\mathbf{u}_j = \mathbf{v}_j$  for  $j = 1, \dots, n$ .

## Definition (Injective and surjective)

Let  $T: U \rightarrow V$ .  $T$  is *injective* if  $T\mathbf{u} = T\mathbf{v}$  implies  $\mathbf{u} = \mathbf{v}$ , if and only if  $\text{null } T = \{\mathbf{0}\}$ .  $T$  is *surjective* if  $\forall \mathbf{v} \in V, \exists \mathbf{u} \in U$  such that  $\mathbf{v} = T\mathbf{u}$ , i.e. if  $\text{range } T = V$ .

## Theorem (Rank Nullity Theorem)

Let  $T: U \rightarrow V$  be a linear transformation, where  $U$  and  $V$  are finite-dimensional vector spaces. Then

$$\text{rank } T + \text{nullity } T = \dim U.$$

$$\dim \text{Im } T + \dim \text{ker } T = \dim U$$

$$U \xrightarrow{S} V \xrightarrow{T} W$$

## Definition (Product of linear maps)

Let  $S \in \mathcal{L}(U, V)$  and  $T \in \mathcal{L}(V, W)$ . We define the product  $ST \in \mathcal{L}(U, W)$  for  $\mathbf{u} \in U$  as  $\underline{ST(\mathbf{u}) = S(T(\mathbf{u}))}$ .

## Definition


A linear map  $T: U \rightarrow V$  is *invertible* if there exists a linear map  $S: V \rightarrow U$  such that  $ST$  is the identity map on  $U$  and  $TS$  is the identity map on  $V$ . Such a map  $S$  is called the *inverse* of  $T$ .

If  $T$  is invertible, we denote the inverse by  $T^{-1}$ . This is justified by the fact that the inverse is unique, as the next proposition shows.

$$V \xrightarrow{S} U \xrightarrow{T} V$$


  
 $\text{Id}_U$

$$V \xrightarrow{T} V \xrightarrow{S} U$$


  
 $ST = \text{Id}_V$

## Proposition

Any invertible linear map has a unique inverse.

*Proof.* Let  $T: V \rightarrow V$  is invertible.

Let  $S$  and  $S'$  be inverses of  $T$ .

$$ST = S'T = \text{Id}_V, \quad TS' = TS = \text{Id}_V$$

For any  $u \in V$ , we will show  $Su = S'u$ .

$$\begin{aligned} Su &= ST(S'u) \quad \text{since } ST = \text{Id}_V \\ &= S(TS'u) \\ &= S(TS'u) \quad \text{since } TS' = \text{Id}_V \end{aligned}$$

$$= S'(S^n)$$

$$= S^n \quad \text{since } S' = \text{Id}_V$$

Since  $S^n = S'^n$  for any  $n \in \mathbb{N}$ , we conclude  $S = S'$ .

## Theorem

A linear map is invertible if and only if it is injective and surjective.

See proof in the book.

## Definition

An invertible linear map is called an *isomorphism*. If there exists an isomorphism from one vector space to another, we say that the vector spaces are *isomorphic*.



## Theorem

Two finite-dimensional vector spaces over  $\mathbb{F}$  are isomorphic if and only if they have the same dimension.

Proof. ( $\Rightarrow$ ) Suppose  $U$  and  $V$  are isomorphic.

$\exists T : U \rightarrow V$  isomorphism (invertible)

By Rank-Nullity theorem

$$\dim U = \dim \text{Im } T + \dim \text{ker } T \quad \dots (*)$$

Since  $T$  is injective and surjective, we have

$$\dim \text{Im } T = \dim V,$$

$$\dim \text{ker } T = 0.$$

Therefore (\*) becomes  $\dim U = \dim V.$



( $\Leftarrow$ ) Suppose  $\dim U = \dim V = n$ .

Then we have a basis  $\{u_i\}_{i=1, \dots, n}$  of  $U$  and  $\{v_i\}_{i=1, \dots, n}$  of  $V$ .

Define  $T: U \rightarrow V$  by

$$T u_i = v_i \quad \text{for } i=1, \dots, n.$$

Then  $T$  is isomorphism  $U \rightarrow V$  (Check!)

# Linear maps and matrices

## Example

Let  $A \in M_{m \times n}$  be a fixed matrix. Then, we can define a linear map  $T_A: \mathbb{F}^n \rightarrow \mathbb{F}^m$  via  $T_A(\mathbf{v}) = A\mathbf{v}$ , where we recall matrix vector multiplication  $(A\mathbf{v})_i = \sum_{k=1}^n A_{ik}v_k$  for  $i = 1, \dots, m$ .

Next we will see that we can use matrices to represent linear maps between finite dimensional vector spaces.

## Definition

Let  $T \in \mathcal{L}(U, V)$  where  $U$  and  $V$  are vector spaces. Let  $\mathbf{u}_1, \dots, \mathbf{u}_n$  and  $\mathbf{v}_1, \dots, \mathbf{v}_m$  be bases for  $U$  and  $V$  respectively. The matrix of  $T$  with respect to these bases is the  $m \times n$  matrix  $\mathcal{M}(T)$  with entries  $A_{ij}$ ,  $i = 1, \dots, m$ ,  $j = 1, \dots, n$  defined by

$$T\mathbf{u}_k = \underbrace{A_{1k}\mathbf{v}_1 + \dots + A_{mk}\mathbf{v}_m}_{eV} \quad \text{unique representation of } T\mathbf{u}_k \text{ by a basis } \{\mathbf{v}_i\}$$

i.e. the  $k$ th column of  $A$  is the scalars needed to write  $T\mathbf{u}_k$  as a linear combination of the basis of  $V$ :

$$T\mathbf{u}_k = \sum_{i=1}^m A_{ik}\mathbf{v}_i \quad \mathcal{M}(T) = (A_{ij})_{\substack{i=1, \dots, m \\ j=1, \dots, n}}$$

Note that since a linear map  $T \in \mathcal{L}(U, V)$  is uniquely determined by its image on a basis of  $U$ , we see that once we pick basis of  $U$  and  $V$  its matrix representation is uniquely determined.

## Example

Polynomials, deg  $\leq 4$ .

Let  $D \in \mathcal{L}(\mathbb{P}_4(\mathbb{R}), \mathbb{P}_3(\mathbb{R}))$  be the differentiation map,  $Dp = p'$ . Find the matrix of  $D$  with respect to the standard bases of  $\mathbb{P}_3(\mathbb{R})$  and  $\mathbb{P}_4(\mathbb{R})$ .

Standard basis:  $1, x, x^2, x^3, (x^4)$   
 $= u_1, u_2, u_3, u_4, u_5$

$$T(u_1) = 0$$

$$T(u_2) = 1 = u_1$$

$$T(u_3) = 2x = 2u_2$$

$$T(u_4) = 3x^2 = 3u_3$$

$$T(u_5) = 4x^3 = 4u_4$$

The matrix is:

$$M(T) = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 2 & 0 & 0 \\ 0 & 0 & 0 & 3 & 0 \\ 0 & 0 & 0 & 0 & 4 \end{pmatrix} \in \mathbb{R}^{4 \times 5}$$



- Observe that if we choose bases  $\mathbf{u}_1, \dots, \mathbf{u}_n$  and  $\mathbf{v}_1, \dots, \mathbf{v}_m$  for  $U, V$  and represent  $T \in \mathcal{L}(U, V)$  as a matrix  $\mathcal{M}(T)$ , then the corresponding map can be obtained by just working with the coordinates of vectors in  $U, V$  with respect to the chosen basis
- If  $\mathbf{u} = \sum_{i=1}^n \alpha_i \mathbf{u}_i$ , then the coordinates of  $T(\mathbf{u})$  with respect to  $\mathbf{v}_1, \dots, \mathbf{v}_m$  can be obtained by the matrix vector multiplication  $\mathcal{M}(T)\boldsymbol{\alpha}$ , where  $\boldsymbol{\alpha}$  is the  $n \times 1$  matrix with entries  $\alpha_j$

## Example

If we want to find the derivative of  $p = x^4 + 12x^3 - 5x^2 + 7$  with respect to the standard monomial basis of  $\mathbb{P}_4(\mathbb{R})$ , we use  $\mathcal{M}(D)$  from the previous example to obtain

$$\mathcal{M}(D)\alpha = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 2 & 0 & 0 \\ 0 & 0 & 0 & 3 & 0 \\ 0 & 0 & 0 & 0 & 4 \end{pmatrix} \begin{pmatrix} 7 \\ 0 \\ -5 \\ 12 \\ 1 \end{pmatrix} = \begin{pmatrix} 0 \\ -10 \\ 36 \\ 4 \end{pmatrix} \begin{matrix} \rightarrow \text{const} \\ \rightarrow x \\ \rightarrow x^2 \\ \rightarrow x^3 \end{matrix}$$

Thus, translating back into the monomial basis of  $\mathbb{P}_3(\mathbb{R})$  gives  
 $D(p) = -10x + 36x^2 + 4x^3$ .

## Other points

- Looking at matrices as representations of linear maps gives us an intuitive explanation for why we do matrix multiplication the way we do! In fact, we want matrix multiplication to represent composition of linear maps
- We can use matrices to solve linear systems.

# Determinants



# Determinant

- The determinant is a function from  $M_{n \times n} \rightarrow \mathbb{F}$ , i.e. it is a function from the entries of a square matrix to a real or complex number.
- Notation:  $\det(A) = |A|$
- The determinant has applications in solving linear systems, computing eigenvalues, etc

## Example: $2 \times 2$ matrix

The determinant of a  $2 \times 2$  matrix is

$$\begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc$$

## Example: $3 \times 3$ matrix

There is a **trick** for finding the determinant of a 3 by 3 matrix:

$$\begin{vmatrix} a & b & c \\ d & e & f \\ g & h & i \end{vmatrix}$$

$$|A| = ace + bfg + cdh - afh - bdi - ceg.$$

# Cofactor expansion

For other  $n \times n$  matrices, one can compute the determinant using **cofactor expansion**.

## Definition (Cofactor expansion)

Let  $A = \{a_{j,k}\}_{j,k=1}^n$  be a  $n \times n$  matrix. Let  $M_{j,k}$  denote the determinant of the  $(n-1) \times (n-1)$  matrix obtained by removing the  $j^{\text{th}}$  row and the  $k^{\text{th}}$  column of  $A$ . For each row  $j = 1, \dots, n$

$$|A| = \sum_{k=1}^n a_{j,k} (-1)^{j+k} M_{j,k}.$$



Similarly, for each column  $k = 1, \dots, n$

$$|A| = \sum_{j=1}^n a_{j,k} (-1)^{j+k} M_{j,k}.$$



The numbers  $C_{j,k} = (-1)^{j+k} M_{j,k}$  are called *cofactors*.

$$A = \begin{pmatrix} & a_{11} & & \\ & \vdots & & \\ c_{j1} & a_{jk} & c_{jn} & \\ & \vdots & & \\ & a_{m1} & & \end{pmatrix} \leftarrow \text{remove}$$

$M_{jk}$  = determinant of the matrix  
obtained by removing  $j$ th row and  $k$ th column of  $A$ .

## Proposition

The determinant of a diagonal matrix or triangular matrix is the product of the entries on the diagonal.

Sketch of proof. Let  $A = \begin{pmatrix} a_{11} & & * \\ 0 & \dots & \\ & & a_{nn} \end{pmatrix}$  upper triangular matrix.  
By induction.

$$n=1 \quad A = (a_{11}) \quad \text{so } \det A = a_{11}.$$

$$\text{Suppose for } n-1 \quad \det A = \prod_{i=1}^{n-1} a_{ii}.$$

For  $n$ . by cofactor expansion

$$A = \begin{pmatrix} a_{11} & & & \\ 0 & & & \\ \vdots & & & \\ 0 & & & \\ & a_{22} & & \\ & & \ddots & \\ & & & a_{nn} \end{pmatrix}$$

use the 1st column.

Since  $a_{11}$  is the only non-zero entry in the 1st column

$$\det A = a_{11} \cdot (-1)^{1+1} \underbrace{M_{11}}_I$$

$$\det \begin{pmatrix} a_{22} & & \\ & \ddots & \\ 0 & & a_{nn} \end{pmatrix} = \prod_{i=2}^n a_{ii}$$

by inductive hypothesis.

$$= a_{11} \cdot \prod_{i=2}^n a_{ii} = \prod_{i=1}^n a_{ii}$$

# Inverse of a matrix

$$C_{j,k} = (-1)^{j+k} M_{j,k}$$

## Theorem

Let  $A$  be an  $n \times n$  invertible matrix and let  $C = \{C_{j,k}\}_{j,k=1}^n$  be its cofactor matrix. Then

$$A^{-1} = \frac{1}{|A|} C^T$$

Note: The matrix  $A$  is invertible if and only if the linear map represented by the matrix is an isomorphism.



# Cramer's rule

## Corollary

Suppose  $A$  is an  $n \times n$  invertible matrix. The linear system  $A\mathbf{x} = \mathbf{b}$  has a unique solution given by

$$x_i = \frac{|A_i|}{|A|}, \quad i = 1, \dots, n,$$

where  $A_i$  is the matrix obtained by replacing the  $i^{\text{th}}$  column of  $A$  with  $\mathbf{b}$ .

$$A = (a_1 \quad \dots \quad a_i \quad \dots \quad a_n)$$

↓ replace.

$$A_i = (a_1 \quad \dots \quad \mathbf{b} \quad \dots \quad a_n)$$

# Transpose of a matrix

## Definition

The *transpose* of an  $m \times n$  matrix  $A$  is the  $n \times m$  matrix, denoted  $A^T$ , defined entry-wise as  $\{A_{j,k}^T\} = \{A_{k,j}\}$  for  $j = 1, \dots, m$  and  $k = 1, \dots, n$  (i.e. the rows of  $A$  are the columns of  $A^T$  and the columns of  $A$  are the rows of  $A^T$ )

$$A = \begin{pmatrix} 1 & 2 \\ 3 & 4 \\ 5 & 6 \end{pmatrix}, \quad A^T = \begin{pmatrix} 1 & 3 & 5 \\ 2 & 4 & 6 \end{pmatrix}$$

# Properties of the determinant

## Proposition

$|A| \neq 0$  if and only if  $A$  is invertible

## Proposition

Let  $A$  be an  $n \times n$  real matrix.

- 1 If  $A$  has a zero column, then  $|A| = 0$ .
- 2 If  $A$  has two equal columns, then  $|A| = 0$ .
- 3 If one column of  $A$  is a multiple of another, then  $|A| = 0$ .
- 4  $|AB| = |A||B|$
- 5  $|\alpha A| = \alpha^n |A|$  for  $\alpha \in \mathbb{F}$
- 6  $|A^T| = |A|$

# Inner product spaces

# Complex numbers

$$a, b \in \mathbb{R}.$$

Recall that for a complex number  $z = a + ib$ , we define the following:

- Real part:  $\operatorname{Re}(z) = a$ ,
- Imaginary part:  $\operatorname{Im}(z) = b$ ,
- Complex conjugate:  $\bar{z} = a - ib$ ,
- Modulus:  $|z| = \sqrt{\operatorname{Re}(z)^2 + \operatorname{Im}(z)^2} = \sqrt{a^2 + b^2}$

We have  $|z|^2 = z\bar{z}$  and  $\operatorname{Re}(z) = \frac{z + \bar{z}}{2}$ .

## Definition

Let  $V$  be an  $\mathbb{F}$ -vector space. A function  $\langle \cdot, \cdot \rangle: V \times V \rightarrow \mathbb{F}$  is called *inner product* on  $V$  if the following holds:

- 1 (Conjugate) symmetry:  $\langle \mathbf{x}, \mathbf{y} \rangle = \overline{\langle \mathbf{y}, \mathbf{x} \rangle}$  for all  $\mathbf{x}, \mathbf{y} \in V$ , where  $\bar{a}$  denotes the complex conjugate for  $a \in \mathbb{C}$
- 2 Linearity in the first argument:  $\langle \alpha \mathbf{x} + \beta \mathbf{y}, \mathbf{z} \rangle = \alpha \langle \mathbf{x}, \mathbf{z} \rangle + \beta \langle \mathbf{y}, \mathbf{z} \rangle$  for all  $\mathbf{x}, \mathbf{y}, \mathbf{z} \in V$  and  $\alpha, \beta \in \mathbb{F}$
- 3 Positive definiteness:  $\langle \mathbf{x}, \mathbf{x} \rangle \geq 0$  and  $\langle \mathbf{x}, \mathbf{x} \rangle = 0$  if and only if  $\mathbf{x} = \mathbf{0}$

A vector space equipped with an inner product is called an *inner product space*.

$$\langle \mathbf{z}, \alpha \mathbf{x} + \beta \mathbf{y} \rangle = \overline{\langle \alpha \mathbf{x} + \beta \mathbf{y}, \mathbf{z} \rangle} = \overline{\alpha \langle \mathbf{x}, \mathbf{z} \rangle + \beta \langle \mathbf{y}, \mathbf{z} \rangle}$$

$$= \bar{\alpha} \overline{\langle \mathbf{x}, \mathbf{z} \rangle} + \bar{\beta} \overline{\langle \mathbf{y}, \mathbf{z} \rangle} = \bar{\alpha} \langle \mathbf{z}, \mathbf{x} \rangle + \bar{\beta} \langle \mathbf{z}, \mathbf{y} \rangle$$

w.r.t. 2nd argument

it's not linear.

## Example

- Standard inner product on  $\mathbb{R}^n$ :  $\langle \mathbf{x}, \mathbf{y} \rangle = \sum_{i=1}^n x_i y_i$  for  $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$
- Standard inner product on  $\mathbb{C}^n$ :  $\langle \mathbf{x}, \mathbf{y} \rangle = \sum_{i=1}^n x_i \bar{y}_i$  for  $\mathbf{x}, \mathbf{y} \in \mathbb{C}^n$   
 $= \mathbf{x}^T \bar{\mathbf{y}}$  ← conjugate.
- On the space of polynomials  $\mathbb{P}_n(\mathbb{R})$ :  $\langle \mathbf{p}, \mathbf{q} \rangle = \int_{-1}^1 p(x)q(x)dx$  for  $\mathbf{p}, \mathbf{q} \in \mathbb{P}_n(\mathbb{R})$   
 $= \mathbf{x}^T \bar{\mathbf{y}}$

check definition vs unit  $\Rightarrow$  Exercise

## Proposition

Let  $V$  be an inner product space. Then  $\mathbf{x} = \mathbf{0}$  if and only if  $\langle \mathbf{x}, \mathbf{y} \rangle = 0$  for all  $\mathbf{y} \in V$ .

*Proof.*  $(\Rightarrow)$  Suppose  $\mathbf{x} = \mathbf{0}$ .

Then by linearity

$$\langle \mathbf{0}, \mathbf{y} \rangle = \langle \mathbf{0} + \mathbf{0}, \mathbf{y} \rangle = \langle \mathbf{0}, \mathbf{y} \rangle + \langle \mathbf{0}, \mathbf{y} \rangle$$

$$\therefore \langle \mathbf{0}, \mathbf{y} \rangle = 0 \quad \text{for } \forall \mathbf{y} \in V.$$

$(\Leftarrow)$  Take  $\mathbf{y} = \mathbf{x}$ .

$$\text{Then } \langle \mathbf{x}, \mathbf{x} \rangle = 0$$

But  $\langle \mathbf{x}, \mathbf{x} \rangle = 0$  iff  $\mathbf{x} = \mathbf{0}$  by definition.

$$\text{Thus } \mathbf{x} = \mathbf{0} //$$



# Cauchy-Schwarz Inequality

## Proposition

Let  $V$  be an inner product space. Then

$$|\langle \mathbf{x}, \mathbf{y} \rangle| \leq \sqrt{\langle \mathbf{x}, \mathbf{x} \rangle} \sqrt{\langle \mathbf{y}, \mathbf{y} \rangle}$$

for all  $\mathbf{x}, \mathbf{y} \in V$ .

Proof. Consider  $\langle tx+iy, tx+iy \rangle$  for  $t \in \mathbb{C}$ .

then by definition

$$\begin{aligned} 0 &\leq \langle tx+iy, tx+iy \rangle \\ &= \langle tx, tx \rangle + \langle tx, iy \rangle + \langle iy, tx \rangle + \langle iy, iy \rangle \\ &= \langle x, x \rangle \underbrace{t\bar{t}}_{=|t|^2} + t\langle x, iy \rangle + \bar{t}\langle iy, x \rangle + \langle y, y \rangle \\ &= \langle x, x \rangle |t|^2 + t\langle x, iy \rangle + \overline{t\langle x, iy \rangle} + \langle y, y \rangle \end{aligned}$$

$$\text{let } f = - \frac{\langle x, y \rangle}{\langle x, x \rangle} \quad \left( \begin{array}{l} \text{You need to consider} \\ \text{case } x=0 \text{ separately} \end{array} \right)$$

$$0 \leq \frac{|\langle x, y \rangle|^2}{\langle x, x \rangle} - \frac{|\langle x, y \rangle|^2}{\langle x, x \rangle} - \frac{|\langle x, y \rangle|^2}{\langle x, x \rangle} + \langle y, y \rangle$$

$$\Rightarrow |\langle x, y \rangle|^2 \leq \langle x, x \rangle \langle y, y \rangle$$

## Proposition

Let  $V$  be an inner product space. Then  $\langle \cdot, \cdot \rangle$  induces a norm on  $V$  via  $\|x\| = \sqrt{\langle x, x \rangle}$  for all  $x \in V$ .

*Proof.* Need to show  $\|\cdot\|$  satisfies the conditions of a norm.

1) Positive definite:

$$\|x\| \geq 0 \quad \text{since} \quad \langle x, x \rangle \geq 0.$$

$$\|x\| = 0 \Leftrightarrow \langle x, x \rangle = 0 \Leftrightarrow x = 0 \quad \text{by definiteness of inner product.}$$

2) Scaling:

$$\begin{aligned} \| \alpha x \| &= \sqrt{\langle \alpha x, \alpha x \rangle} = \sqrt{\alpha \bar{\alpha} \langle x, x \rangle} \\ &= \sqrt{|\alpha|^2 \langle x, x \rangle} \end{aligned}$$

$$= |\alpha| \sqrt{\langle x, x \rangle}$$

$$= |\alpha| \|x\|$$

3.) Triangle inequality

$$(\|x\| + \|y\|)^2 - \|x+y\|^2$$

$$= (\|x\|^2 + 2\|x\| \cdot \|y\| + \|y\|^2) - \langle x+y, x+y \rangle$$

$$= \left( \quad \quad \quad \right) - \left( \underbrace{\langle x, x \rangle}_{=\|x\|^2} + \langle x, y \rangle + \underbrace{\langle y, x \rangle}_{=\langle x, y \rangle} + \underbrace{\langle y, y \rangle}_{=\|y\|^2} \right)$$

$$= (\cancel{\|x\|^2} + 2\|x\| \cdot \|y\| + \cancel{\|y\|^2}) - (\cancel{\|x\|^2} + 2\operatorname{Re}\langle x, y \rangle + \cancel{\|y\|^2})$$

$$= 2 \left( \|x\| \cdot \|y\| - \operatorname{Re}\langle x, y \rangle \right) = (*)$$

By C-S,

$$\operatorname{Re}\langle x, y \rangle \leq |\langle x, y \rangle| \stackrel{\text{C-S}}{\leq} \|x\| \cdot \|y\|.$$

Thus, we conclude  $(*) \geq 0$ .



Note: With this identification the Cauchy-Schwarz inequality can be restated as:  
 $|\langle \mathbf{x}, \mathbf{y} \rangle| \leq \|\mathbf{x}\| \|\mathbf{y}\|$  for all  $\mathbf{x}, \mathbf{y} \in V$ .

### Example

The norm introduced by the standard inner product on  $\mathbb{R}^n$  is the Euclidean distance.

# References

Axler S. *Linear Algebra Done Right*. 3rd ed. Undergraduate Texts in Mathematics. Springer, 2015. Available from:  
<https://link.springer.com/book/10.1007/978-3-319-11080-6>

Treil S. *Linear Algebra Done Wrong*. 2017. Available from:  
<https://www.math.brown.edu/streil/papers/LADW/LADW.html>