

Module 5: Metric spaces III

Operational math bootcamp



Statistical Sciences
UNIVERSITY OF TORONTO

Ichiro Hashimoto

University of Toronto

July 15, 2025

Last time

Looked at open and closed sets.

Discussed sequences, which includes Cauchy sequences and subsequences:

- Convergent sequence: $x_n \rightarrow x \Leftrightarrow \forall \epsilon > 0 \exists n_\epsilon \in \mathbb{N}$ s.t. $d(x_n, x) < \epsilon$ for all $n \geq n_\epsilon$
- Cauchy sequence: $\forall \epsilon > 0 \exists n_\epsilon \in \mathbb{N}$ s.t. $d(x_n, x_m) < \epsilon$ for all $n, m \geq n_\epsilon$
- Proved that a convergent sequence is Cauchy
- Discussed complete metric spaces, where all Cauchy sequences converge (like \mathbb{R} with the usual metric, absolute value)
- Proved that a sequence converges to x if and only if all subsequences converge to x

Outline for today

- Continuity
- Equivalent metrics
- Density and separability

Continuity

Definition

Let (X, d_X) and (Y, d_Y) be metric spaces, let $x_0 \in X$, and let $f : X \rightarrow Y$. f is *continuous* at x_0 if for every sequence $(x_n)_{n \in \mathbb{N}}$ in X that converges to x_0 , we have $\lim_{n \rightarrow \infty} f(x_n) = f(x_0)$.

We say that f is continuous if it is continuous at every point in X .

Theorem

Let (X, d_X) and (Y, d_Y) be metric spaces, let $x_0 \in X$, and let $f : X \rightarrow Y$. The following are equivalent:

- (i) f is continuous at x_0
- (ii) for all $\epsilon > 0$, there exists $\delta > 0$ such that $d_Y(f(x), f(x_0)) < \epsilon$ for all $x \in X$ with $d_X(x, x_0) < \delta$
- (iii) for each $\epsilon > 0$, there is $\delta > 0$ such that $B_\delta(x_0) \subseteq f^{-1}(B_\epsilon(f(x_0)))$

- (i) f is continuous at x_0
- (ii) for all $\epsilon > 0$, there exists $\delta > 0$ such that $d_Y(f(x), f(x_0)) < \epsilon$ for all $x \in X$ with $d_X(x, x_0) < \delta$
- (iii) for each $\epsilon > 0$, there is $\delta > 0$ such that $B_\delta(x_0) \subseteq f^{-1}(B_\epsilon(f(x_0)))$

Proof. (i) \Rightarrow (ii)

$$(ii) \Rightarrow (iii)$$

$$(iii) \Rightarrow (i)$$

Corollary

Let (X, d_X) and (Y, d_Y) be metric spaces and let $f : X \rightarrow Y$. The following are equivalent:

- (i) f is continuous
- (ii) if $U \subseteq Y$ is open, then $f^{-1}(U)$ is open
- (iii) if $F \subseteq Y$ is closed, then $f^{-1}(F)$ is closed

We need the following results about sets and functions:

Let X and Y be sets and $f : X \rightarrow Y$. Let $A, B \subseteq Y$. Then

$$\textcircled{1} \quad A \subseteq B \implies f^{-1}(A) \subseteq f^{-1}(B)$$

$$\textcircled{2} \quad f^{-1}(Y \setminus A) = X \setminus f^{-1}(A)$$

Proof. Let (X, d_X) and (Y, d_Y) be metric spaces and let $f : X \rightarrow Y$.

(i) \Rightarrow (ii):

$$(ii) \Rightarrow (i)$$

$$(ii) \Rightarrow (iii)$$

$$(iii) \Rightarrow (ii)$$

Definition

Let (X, d_X) and (Y, d_Y) be metric spaces and let $f : X \rightarrow Y$.

- f is *uniformly continuous* if for all $\epsilon > 0$, there exists $\delta > 0$ such that for every $x_1, x_2 \in X$ with $d_X(x_1, x_2) < \delta$, we have $d_Y(f(x_1), f(x_2)) < \epsilon$
- f is *Lipschitz continuous* if there exists a $K > 0$ such that for every $x_1, x_2 \in X$ we have $d_Y(f(x_1), f(x_2)) \leq K d_X(x_1, x_2)$

Proposition

Let (X, d_X) and (Y, d_Y) be metric spaces and let $f : X \rightarrow Y$.

f is Lipschitz continuous $\Rightarrow f$ is uniformly continuous $\Rightarrow f$ is continuous

Proof is one of your exercises.

Contraction Mapping Theorem

Definition

Let (X, d) be a metric space and let $f : X \rightarrow X$. We say that $x^* \in X$ is a *fixed point* of f if $f(x^*) = x^*$.

Definition

Let (X, d) be a metric space and let $f : X \rightarrow X$. f is a *contraction* if there exists a constant $k \in [0, 1)$ such that for all $x, y \in X$, $d(f(x), f(y)) \leq kd(x, y)$.

Observe that a function is a contraction if and only if it is Lipschitz continuous with constant $K < 1$.

Theorem (Contraction Mapping Theorem)

Suppose that $f : X \rightarrow X$ is a contraction and the metric space X is complete. Then f has a unique fixed point x^ .*

Example

Let $f : \left[-\frac{1}{3}, \frac{1}{3}\right] \rightarrow \left[-\frac{1}{3}, \frac{1}{3}\right]$ be defined by the mapping $x \mapsto x^2$. Assume we use the standard Euclidean metric, $d(x, y) = |x - y|$. f has a unique fixed point because

Equivalent metrics

Definition (Equivalent metrics)

Two metrics d_1 and d_2 on a set X are *equivalent* if the identity maps from (X, d_1) to (X, d_2) and from (X, d_2) to (X, d_1) are continuous.

Proposition

Two metrics d_1, d_2 on a set X are equivalent if and only if they have the same open sets or the same closed sets.

Definition

Two metrics d_1 and d_2 on a set X are *strongly equivalent* if for every $x, y \in X$, there exists constants $\alpha > 0$ and $\beta > 0$ such

$$\alpha d_1(x, y) \leq d_2(x, y) \leq \beta d_1(x, y).$$

If two metrics are strongly equivalent then they are equivalent. The proof of this is one of the exercises.

Example

We show that the Euclidean distance (induced by 2-norm) and the metric induced by the ∞ -norm are equivalent on \mathbb{R}^n .

Can you think of an example that we've seen of a metric that isn't equivalent to the Euclidean metric?



Density

Definition

Let (X, d) be a metric space. A subset $A \subseteq X$ is called *dense* if $\overline{A} = X$.

Why are dense sets important?

Examples

1. $(\mathbb{R}, |\cdot|)$

2. Let X be a set and define $d: X \times X \rightarrow \mathbb{R}$ by

$$d(x, y) = \begin{cases} 0, & x = y, \\ 1, & x \neq y. \end{cases}$$

Definition

A metric space (X, d) is *separable* if it contains a countable dense subset.

Example:

Example

Define $\ell_\infty = \{(x_n)_{n \in \mathbb{N}} : x_n \in \mathbb{R}, \sup_{n \in \mathbb{N}} |x_n| < \infty\}$, the space of bounded real valued sequences. Endow ℓ_∞ with a metric induced by the supremum norm, namely $d((x_n)_{n \in \mathbb{N}}, (y_n)_{n \in \mathbb{N}}) = \sup_{n \in \mathbb{N}} |x_n - y_n|$. Then ℓ_∞ is **not separable** with respect to the topology induced by this metric.

Note: this proof relies on content from the cardinality section. Specifically, Cantor's theorem gives us that $|A| < |\mathcal{P}(A)|$ for any set A .

Proof.

Proof continued.

References

Jiří Lebl (2022). *Basic Analysis I*. Vol. 1. Introduction to Real Analysis.
<https://www.jirka.org/ra/realanal.pdf>

Runde, Volker (2005). *A Taste of Topology*. Universitext. url:
<https://link.springer.com/book/10.1007/0-387-28387-0>